

Microstegium vimineum


Nepalese browntop
Japanese stiltgrass

MIVI

8303



Chris Evans, The University of Georgia, www.forestryimages.org



Voucher specimens needed in ND, SD, NE, KS, MO, IA, MN, WI and MI.

Microstegium vimineum

Nepalese browntop	MIVI	8303
Japanese stiltgrass		

Form: Sprawling annual grass, 0.2 to 1 m tall.

Leaves: Alternate; narrowly lanceolate 5-10 cm long; blades flat and sparsely hairy on both surfaces, as well as along the margins.

Flowers/fruit: Inflorescence a digitate cluster of 2-4 narrow spikelike branches; pubescent; awns sometimes present on lemmas. Spikelets paired (one sessile, one pedicellate); lower glume flat, upper keeled.

Habitat: Floodplains, streamsides, forest edge, roadsides, swamps, wet fields and trailsides.

Notes

Microstegium vimineum
Nepalese browntop, Japanese stiltgrass



Elizabeth J. Czarapata



UGA0016156

May.

James H. Miller, USDA Forest Service,
www.forestryimages.org



Elizabeth J. Czarapata



UGA2307199

Spring.

Ted Bodner, Southern Weed Science
Society, www.forestryimages.org



September.

Chuck Bargeron, The University of
Georgia, www.forestryimages.org



UGA0016155

September.

Ted Bodner, Southern Weed Science Society, www.forestryimages.org



UGA0581028

September.

Jil M. Swearingen, USDI National Park Service, www.forestryimages.org



UGA0016160

December.

James H. Miller, USDA Forest Service, www.forestryimages.org