

Comparative Impact of *Scymnus ningshanensis* and *Pseudoscymnus tsugae* (Coleoptera: Coccinellidae) on the Hemlock Woolly Adelgid

Elizabeth Butin¹, Joseph Elkinton¹, Nathan Havill², and Michael Montgomery²

¹Department of Entomology, University of Massachusetts, Amherst, MA 01003

²USDA Forest Service, Northeastern Center for Forest Health Research,
51 Mill Pond Rd., Hamden, CT 06514

Abstract

The hemlock woolly adelgid (*Adelges tsugae* Annand) is an introduced pest thought to be native to Asia. Damage to eastern hemlock and Carolina hemlock can be serious (Salom et al. 1996), but western and Asian hemlocks are seldom damaged. Potential biological control agents have been observed in Japan and China (Sasaji and McClure 1997, Yu et al. 2000). We compared two of these which have been previously imported, *Scymnus ningshanensis* Yu et Yao, a coccinellid from China (Yu et al. 2000) and *Pseudoscymnus tsugae* Sasaji & McClure a coccinellid from Japan (Sasaji and McClure 1997).

For each lady beetle, we examined the host range and the numerical response to prey density (adelgid egg masses) in the laboratory, and in field studies, their ability to reduce hemlock woolly adelgid population growth. When given a choice between two prey species in the laboratory, *S. ningshanensis* preferred *A. tsugae* to *Adelges laricis* and *Prociphilus tessellatus* ($p < 0.05$, 2 sample t-test) but preferences between *A. tsugae* and *Adelges cooleyi* or *Pineus strobi* were not different ($p > 0.05$, 2 sample t-test). The host range results for *P. tsugae* were inconclusive because the beetles fed very little on all prey species provided, including the hemlock woolly adelgid. *S. ningshanensis* showed a positive numerical response ($p < 0.05$, linear regression), and *P. tsugae* showed a density independent response ($p > 0.05$, linear regression) to an increasing density of hemlock woolly adelgid egg masses. In the field, caged branches with a pair of *S. ningshanensis* resulted in a negative population growth of *A. tsugae*, while cages with a pair of *P. tsugae* adults and the control without lady beetles resulted in an increase in the population growth of *A. tsugae* ($p < 0.05$, ANOVA).

Literature Cited

- Salom, S.M.; Tigner, T.C.; Reardon, R.C. 1996. **Proceedings of the first hemlock woolly adelgid review.** Oct. 12, 1995, Charlottesville, Virginia. USDA, Forest Service, FHTET 96-10.
- Sasaji, H.; McClure, M.S. 1997. **Description and distribution of *Pseudoscymnus tsugae* sp. nov. (Coleoptera: Coccinellidae), an important predator of hemlock woolly adelgid in Japan.** Annals of the Entomological Society of America. 90: 563-568.
- Yu, G.; Montgomery, M.E.; Defu, Y. 2000. **Lady beetles (Coleoptera: Coccinellidae) from Chinese hemlocks infested with the hemlock woolly adelgid, *Adelges tsugae* Annand (Homoptera: Adelgidae).** The Coleopterists Bulletin. 54(2): 154-199.