



United States
Department of
Agriculture

Forest Service

**Northern
Research Station**

General Technical
Report NRS-54




Urban and Community Forests of the North Central East Region

**Illinois
Indiana
Michigan**

**Ohio
Wisconsin**

**David J. Nowak
Eric J. Greenfield**



Abstract

This report details how land cover and urbanization vary within the states of Illinois, Indiana, Michigan, Ohio, and Wisconsin by community (incorporated and census designated places), county subdivision, and county. Specifically this report provides critical urban and community forestry information for each state including human population characteristics and trends, changes in urban and community lands, tree canopy and impervious surface cover characteristics, distribution of land-cover classes, a relative comparison of urban and community forests among local government types, determination of priority areas for tree planting, and a summary of urban tree benefits. Report information can improve the understanding, management, and planning of urban and community forests. The data from this report is reported for each state on the CD provided in the back of this book, and it may be accessed by state at: <http://www.nrs.fs.fed.us/data/urban>.

The Authors

DAVID J. NOWAK is a research forester and project leader, and ERIC J. GREENFIELD is a forester with the Forest Service's Northern Research Station at Syracuse, NY.

Manuscript received for publication September 2009

Published by:
U.S. FOREST SERVICE
11 CAMPUS BLVD SUITE 200
NEWTOWN SQUARE, PA 19073

January 2010

For additional copies:
U.S. Forest Service
Publications Distribution
359 Main Road
Delaware, OH 43015
Fax: (740) 368-0152

Visit our homepage at: <http://www.nrs.fs.fed.us/>



Printed on Recycled Paper

CONTENTS

Introduction	1
Report Overview	2
Urban Forest Attributes	2
Summary	10
Acknowledgments	10
Illinois' Urban and Community Forests	11
Indiana's Urban and Community Forests	19
Michigan's Urban and Community Forests	27
Ohio's Urban and Community Forests	35
Wisconsin's Urban and Community Forests	43
Literature Cited	51
Appendix	54

INTRODUCTION

As part of the Forest and Rangeland Renewable Resources Planning Act of 1974, the first national assessment of urban forests was completed in 2000 (Dwyer et al. 2000, Nowak et al. 2001b). This assessment used 1-km resolution Advanced Very-High Resolution Radiometer (AVHRR) data (Zhu 1994) and 1990 U.S. Census Bureau (2007) population and geographic data to assess urban tree cover. The assessment concluded that urban areas in the conterminous United States doubled in size between 1969 and 1994 and covered 3.5 percent of the total land area. Urban areas were estimated to contain approximately 3.8 billion trees with an average tree canopy cover of 27 percent.

To update this first report, higher resolution (30 m) tree canopy and impervious surface cover maps were used (from 2001 Landsat satellite imagery and published in 2007) (Homer et al. 2007, U.S. Geol. Surv. 2007) in conjunction with 1990 and 2000 census and geographic data (1:5,000,000 scale cartographic boundary files) (U.S. Census Bureau 2007) to assess current urban and community forest attributes. These results are being published for each of the lower 48 United States to provide information on urban change and state-specific urban and community forestry data.

This report includes information for the following states: Illinois, Indiana, Michigan, Ohio, and Wisconsin.

Data are reported for the state, county, county subdivision, and community jurisdictions. The jurisdictional units used in this report are derived from U.S. Census (2007) geographic data and defined legal or statistical divisions. “County”¹ refers to the primary subdivision within states. “County subdivisions” are primary divisions of a county and are statistically equivalent entities for the reporting of census data. They include census county divisions (CCD), census sub areas, minor civil divisions (MCD), and unorganized territories. “Communities” are incorporated and census designated places, and consolidated cities (U.S. Census Bureau 2007). For detailed definitions, see http://www.census.gov/geo/www/cob/cs_metadata.html (2007).

¹The primary legal divisions of most states are termed “counties.” In Louisiana, these divisions are known as “parishes.” In four states (Maryland, Missouri, Nevada, and Virginia), there are one or more incorporated places that are independent of any county organization and thus constitute primary divisions of their states; these incorporated places are known as “independent cities” and are treated as equivalent to counties for statistical purposes. (For some statistical purposes they may be treated as county subdivisions and places.) The District of Columbia has no primary divisions, and the entire area is considered equivalent to a county for statistical purposes. (http://www.census.gov/geo/www/cob/co_metadata.html, 2007)

REPORT OVERVIEW

The information in this report can aid local and regional managers and planners of urban and community forest resources. This report provides urban and community forest reference information and data from the state to local level on the following attributes related to the urban and community forest resource:

- Human population characteristics and trends
- Urban and community land
- Tree canopy cover characteristics
- Impervious surface cover characteristics
- Classified land-cover characteristics
- Relative comparisons of urban and community forests
- Priority areas for tree planting
- Urban tree benefits

Information in this report can be used by urban and community forestry professionals to:

- Understand general land-cover characteristics and urbanization trends at several geographic scales
- Compare tree canopy cover among similar communities
- Determine areas of greatest growth and areas of highest tree planting priority
- Relate urban and community forests to pollution removal and carbon storage
- Promote more detailed and/or locally appropriate urban and community forest inventories, censuses, or field surveys (e.g., i-Tree – www.itreetools.org)
- Establish local to statewide standards related to urban and community forestry
- Support urban and community forestry programs
- Improve urban and community forest management and planning

The remainder of this section details how information was derived for each attribute reported for the urban and community areas. The subsequent state summaries

detail the findings for each state in this region. Most tables for each state are not given in this report, rather they can be found on the CD provided with this report or accessed at: <http://www.nrs.fs.fed.us/data/urban>.

URBAN FOREST ATTRIBUTES

Human Population Characteristics and Trends

Human population and population density changes over time, and geographic distribution are important measurements of the urban environment because human populations are an integral part of community and urban forest dynamics. Within divisions of state, county, county subdivision, and community, total population, population changes from 1990 to 2000 and population density are detailed based on U.S. Census data (U.S. Census Bureau 2007).

Urban and Community Land

Two geographic definitions overlap: “community” and “urban”. The definition of community is based on jurisdictional or political boundaries delimited by U.S. Census definitions of places (U.S. Census Bureau 2007). Community lands are places of established human settlement that may include all, some, or no urban land within their boundaries.

The definition of urban is based on population density as delimited using the U.S. Census Bureau’s (2007) definition: all territory, population, and housing units located within urbanized areas or urban clusters. Urbanized area and urban cluster boundaries encompass densely settled territories, which are described by one of the following:

- One or more block groups or census blocks with a population density of at least 386.1 people/km² (1,000 people/mile²)
- Surrounding block groups and census blocks with a population density of 193.1 people/km² (500 people/mile²)
- Less densely settled blocks that form enclaves or indentations, or are used to connect discontinuous areas

More specifically, urbanized areas consist of territory with 50,000 or more people. Urban clusters, a concept new to the 2000 Census, consist of territory with at least 2,500 people but fewer than 50,000 people. This new definition tends to be more restrictive than the 1990 U.S. Census urban definition and encompasses many areas typically considered suburban. The 2000 Census definition of urban was applied to 1990 Census geographic data to analyze change in urban land between 1990 and 2000 (Nowak et al. 2005).

As urban land reveals the more heavily populated areas (population density-based definition) and community land indicates both urban and rural (i.e., non-urban) communities that are recognized by their geopolitical boundaries (political definition), both definitions provide information related to human settlements and the forest resources within those settlements. As some urban land exists beyond community boundaries and not all community land is urban (i.e., communities are often a mix of urban and rural land), the category of “urban or community” was created to understand forest attributes accumulated by the union of these two definitions. The “urban or community” term used throughout this report encompasses both urban land and land in communities.

Percent urban land is a ratio of urban land over total land within a census geographic division, and percent community land is a ratio of community land over total land within the geopolitical unit. In addition, changes in urban land and changes in community land are reported between 1990 and 2000.

For each state, Tables 1 through 4 summarize the population, and urban and community land attributes for the state, communities, county subdivisions, and counties respectively (CD and <http://www.nrs.fs.fed.us/data/urban>).

Tree Canopy Cover Characteristics

Tree canopy cover is a critical measure of the urban and community forest resource. Tree canopy cover gives a broad indication of the overall forest resource and its associated benefits. To assess urban and

community land cover characteristics, the multi-resolution land characteristics consortium’s National Land Cover Database (NLCD) was used (Homer et al. 2004, U.S. Geol. Surv. 2007, Yang et al. 2003). The NLCD, released in early 2007, was processed from 2001 Landsat satellite imagery and provides estimates of percentage tree canopy and impervious surface cover within 30-m pixels or cells across the state. The tree canopy percentages in this report are calculated using the land area (not including water) of the geopolitical units derived from the U.S. Census cartographic boundary data and NLCD. In addition to percentage tree cover, four other canopy cover attributes, derived from the same data, were assessed:

- Tree canopy cover per capita—Tree canopy cover (m²) divided by the number of people within the area of analysis.
- Total green space—Total area minus impervious and water cover (ha). This attribute estimates pervious cover (i.e., grass, soil, or tree-covered areas).
- Canopy green space—Tree cover divided by total green space (percent). This value is the proportion of the total green space that is filled by tree canopies.
- Available green space—Total green space minus tree canopy cover (ha). This value is the amount of grass and soil area not covered with tree canopies and potentially available for planting.

Impervious Surface Cover Characteristics

Similar to tree cover, impervious surface cover provides another piece of valuable information related to the urban environment. Impervious surface cover gives an indication of an area’s developed hardscape, which has important influences on urban air temperatures and water flows and also yields information on limitations to urban tree cover. Impervious surface cover also was derived from the NLCD database (U.S. Geol. Surv. 2007). The impervious surface cover percentages in this report are calculated using the land area (not including water) of the geopolitical units derived from the U.S. Census

cartographic boundary data and NLCD. Impervious surface per capita is calculated from NLCD 2001 and U.S. Census data.

For each state, Tables 1, and 5 through 7 summarize the tree canopy and impervious surface cover attributes for the state, communities, county subdivisions, and counties respectively (CD and <http://www.nrs.fs.fed.us/data/urban>).

Classified Land-cover Characteristics

Land-cover types also are summarized using 2001 Landsat satellite data that were classified with the U.S. Geological Survey land cover categorization scheme based on a modified Anderson land-cover classification (U.S. Geol. Surv. 2007). Land area, tree canopy cover, and available green space within generalized land cover categories vary among communities, county subdivisions, counties, and state. The percentages are calculated from the NLCD 2001 and U.S. Census cartographic boundary data. The land-cover categories defined here are derived from established NLCD 2001 land-cover classes. These generalized land-cover categories or types may not be present in some states.

- Developed—NLCD classes 21 (developed-open space), 22 (developed-low intensity), 23 (developed-medium intensity), and 24 (developed-high intensity)
- Barren—NLCD class 31 (barren land [rock/sand/clay])
- Forested—NLCD classes 41 (deciduous forest), 42 (evergreen forest), and 43 (mixed forest)
- Shrub/Scrub—NLCD class 52 (shrub/scrub)
- Grassland—NLCD class 71 (grassland/herbaceous)
- Agriculture—NLCD classes 81 (pasture/hay) and 82 (cultivated crops)
- Wetland—NLCD classes 90 (woody wetlands) and 95 (emergent herbaceous wetlands)

For each state, Tables 8 through 10 summarize the classified land-cover characteristics for communities, county subdivisions, and counties and state respectively (CD and <http://www.nrs.fs.fed.us/data/urban>).

Relative Comparisons of Tree Cover

A question commonly asked in evaluating the urban and community forest resource is, “How does my community compare with other communities?”

To help answer this question, tree canopy cover was compared among the counties, county subdivisions, and communities relative to other areas with comparable population density and within the same NLCD mapping unit (ecoregion). For this comparison, seven population density classes were established:

- Density class 1 — 0 to 38.6 people/km² (0 to 99.9 people/mile²)
- Density class 2 — 38.7 to 96.5 people/km² (100 to 249.9 people/mile²)
- Density class 3 — 96.6 to 193.1 people/km² (250 to 499.9 people/mile²)
- Density class 4 — 193.2 to 289.6 people/km² (500 to 749.9 people/mile²)
- Density class 5 — 289.7 to 386.2 people/km² (750 to 999.9 people/mile²)
- Density class 6 — 386.3 to 1931.2 people/km² (1000 to 4999.9 people/mile²) and
- Density class 7 — 1931.3 or greater people/km² (5000 or greater people/mile²)

Mapping zones were delimited within the NLCD to increase classification accuracy and efficiency (Fig. A). The mapping units represent relatively homogeneous ecological conditions (Homer and Gallant 2001). To locate geopolitical units within a mapping zone, centroid (geometric center) points of the local governments were used.

For three or more geographic units in the same mapping zone and population density class, a standardized tree canopy score based on the range of values within that zone and class was assigned to each unit. The standardized score is calculated as:

Standardized score = (tree canopy percent of unit – minimum tree canopy percentage in class)/range of tree canopy percent in class.



Figure A.—The mapping zones of the continental United States relative to states and land cover (NLCD 2001).

Communities, county subdivisions, and counties were assigned to one of the following categories based on their standardized score:

- Excellent—Standardized score of 0.9 to 1.0
- Very Good—0.7 to 0.89
- Good—0.5 to 0.69
- Fair—0.3 to 0.49
- Poor—0 to 0.29

To help understand the variability of tree cover, minimum, median, maximum, and weighted mean values for percent tree canopy cover in each population density class of each political subdivision are reported in Table 11 for each mapping zone (CD and <http://www.nrs.fs.fed.us/data/urban>). This information can be used to understand the actual range and values used for the assessment.

For each state, Tables 12 through 14 summarize the urban and community forest ratings for communities, county subdivisions, and counties respectively (CD and <http://www.nrs.fs.fed.us/data/urban>).

Priority Areas for Tree Planting

NLCD (U.S. Geol. Survey 2007) and 2000 U.S. Census data (2007) were used to produce an index that prioritizes tree planting areas for communities, county subdivisions, and counties. An index was developed to help identify areas with relatively low tree canopy cover and high population density (high priority tree-planting areas). This index provides one form of prioritization. States and local governments may design their own prioritization method incorporating individual and diverse value systems. The index used in this report combines three criteria.

- Population density—The greater the population density, the greater the priority for tree planting
- Canopy green space—The lower the value, the greater the priority for tree planting
- Tree canopy cover per capita—The lower the amount of tree canopy cover per person, the greater the priority for tree planting

Each criterion above was standardized² on a scale of 0 to 1, with 1 representing the maximum population density and minimum canopy green space and tree cover per capita. The standardized values were weighted to produce a combined score:

$$I = (PD * 40) + (CG * 30) + (TPC * 30)$$

Where I is the combined index score

PD is the standardized population density value

CG is the standardized canopy green space value, and

TPC is the standardized tree cover per capita value.

The combined score was standardized again and multiplied by 100 to produce the planting priority index. The tree planting priority index (PPI) ranks each state's communities, county subdivisions, and counties with values from 100 (highest priority) to 0 (lowest priority). This index is a type of "environmental equity" index with areas of higher human population density and lower canopy green space and tree cover per capita tending to get the higher index value.

For each state, Tables 15 through 17 summarize the tree planting priority index for communities, county subdivisions, and counties respectively (CD and <http://www.nrs.fs.fed.us/data/urban>).

Urban Tree Benefits

Urban and community forests are important for human and ecological health (Nowak and Dwyer 2007). The benefits ascribed to urban and community trees include:

- Carbon storage and sequestration
- Air pollution removal
- Surface air temperature reduction
- Reduced building energy use
- Absorption of ultraviolet radiation
- Improved water quality
- Reduced noise pollution
- Improved human comfort
- Increased property value
- Improved human physiological and psychological well-being
- Improved aesthetics
- Improved community cohesion

To understand the contribution and magnitude of the forest resource in urban or community areas, the total number of trees, carbon storage and annual carbon uptake (sequestration), air pollution removal, and the associated dollar values for carbon and air pollution benefits are estimated.

Carbon sequestration and storage values were estimated from tree cover (m²) multiplied by average carbon storage (9.1 kg C/m²), and sequestration (0.3 kg C/m²) density values derived from several U.S. communities (e.g., Nowak and Crane 2002). Monetary values associated with urban tree carbon storage and sequestration were based on the 2001-2010 projected marginal social cost of carbon dioxide emissions, \$22.8/t C (Fankhauser 1994). The number of urban and community trees was estimated in a similar manner multiplying tree canopy cover (m²) by average tree density per hectare of canopy cover from several U.S. cities (Table A).

Air pollution removal estimates are derived from the Urban Forest Effects (UFORE) model (Nowak and Crane 2000) and 2000 weather and pollution data (National Climatic Data Center 2000, U.S. EPA 2008). The UFORE model was used to integrate hourly pollution and weather data with urban or community tree cover data to estimate annual pollution removal in each state (Nowak and Crane 2000, Nowak et al. 2006d).

²Standardized value for population density (PD) was calculated as $PD = (n - \min) / r$, where PD is the value 0-1, n is the value for the geopolitical unit (population/km²), min is the minimum value for all units, and r is the range of values among all units (maximum value – minimum value). Standardized value for canopy green space (CG) was calculated as $CG = (\max - n) / r$, where CG is the value 0-1, max is the maximum value for all geopolitical units, n is the value for the unit (tree canopy cover m²/total green space m²), and r is the range of values. Standardized value for tree cover per capita (TPC) was calculated as $TPC = (\max - n) / r$, where TPC is the value (0-1), max is the maximum value for all geographic units, n is the value for the geopolitical unit (m²/capita), and r is the range of values among all units.

Table A.—Average number of trees, carbon storage, and carbon sequestration rates per unit of canopy cover for several U.S. cities

City	Trees (no./ha cover)	Carbon	
		Storage (kg C/m ² cover)	Sequestration (kg C/m ² cover)
Atlanta, GA ^a	751.5	9.7	0.3
Baltimore, MD ^a	598.1	12.3	0.3
Boston, MA ^a	371.7	9.1	0.3
Chicago, IL ^b	618.0	12.9	n/a
Casper, WY ^c	252.8	7.0	0.2
Freehold, NJ ^a	275.0	10.4	0.3
Jersey City, NJ ^a	308.7	4.4	0.2
Minneapolis, MN ^d	245.5	5.7	0.2
Moorestown, NJ ^a	547.9	9.9	0.3
Morgantown, WV ^a	829.6	10.6	0.3
New York, NY ^e	312.0	7.3	0.2
Philadelphia, PA ^f	394.3	9.0	0.3
San Francisco, CA ^g	468.1	12.3	0.3
Syracuse, NY ^h	583.1	10.5	0.3
Oakland, CA ⁱ	570.0	5.2	n/a
Washington, DC ^j	423.4	10.4	0.3
Woodbridge, NJ ^a	557.3	8.2	0.3
Mean	476.9	9.1	0.3

^a Unpublished data analyzed using UFORE model

^b Nowak 1994a,b

^c Nowak et al. 2006a

^d Nowak et al. 2006b

^e Nowak et al. 2007a

^f Nowak et al. 2007b

^g Nowak et al. 2007c

^h Nowak et al. 2001a

ⁱ Nowak 1993; Nowak and Crane 2002

^j Nowak et al. 2006c

To estimate pollution by urban trees in each state, state pollutant flux rates (grams of pollution removal per square meter of canopy per year) were derived from a study of national pollution removal by urban trees for the year 1994 (Nowak et al. 2006d). As pollution concentrations vary through time, the 1994 flux rates were adjusted to 2000 values based on average regional pollution concentration changes between 1994 and 2000 (U.S. EPA 2003). As

flux rate = deposition velocity * pollution concentration,

the ratio of the pollution concentration between years was used to update the flux rate. Arithmetic mean concentration values were used for nitrogen dioxide, particulate matter less than 10 microns, and sulfur dioxide, 2nd Max. 8-hr average for carbon dioxide, and 4th Max. 8-hr average for ozone, to determine

the ratio of change between 1994 and 2000 (U.S. EPA 2003). The new 2000 flux rates were multiplied by urban or community tree cover in the state to estimate total pollution removal by trees.

Pollution removal dollar value estimates were calculated using 1994 national median externality values used in energy decision making (Murray et al. 1994, Ottinger et al. 1990). The 1994 values were adjusted to 2007 dollars based on the producer price index (U.S. Dept. of Labor 2008). These values, in dollars/metric ton (t) are:

- Nitrogen dioxide (NO₂) = \$9,906/t
- Particulate matter less than 10 microns (PM₁₀) = \$6,614/t
- Sulfur dioxide (SO₂) = \$2,425/t
- Carbon monoxide (CO) = \$1,407/t

Externality values for ozone (O₃) were set to equal the value for NO₂. Externality values can be considered the estimated cost of pollution to society that is not accounted for in the market price of the goods or services that produced the pollution.

For each state, Table 1 summarizes carbon storage and air pollution removal estimates for urban, community, and urban or community trees statewide.

Data Accuracy and Application

The data presented in this report yield the most comprehensive and up-to-date assessment of continental U.S. urban and community forests. The data allows for relative comparisons among geographies and provides baseline information for assessing relative changes in urban and community forest cover in the future. As stated previously, tree cover information was based on finer resolution data than used in the original urban forest assessment (Dwyer et al. 2000). As the methodologies for quantifying tree cover have changed between the original and current assessment, evaluating changes is not possible since the detected changes could be caused by either actual landscape changes or differences in methodology.

The U.S. Census generalized cartographic boundary data are a simplified and smoothed extracts of the Topologically Integrated Geographic Encoding and Referencing (TIGER) database, with a target scale range of 1:5,000,000 (U.S. Census Bureau 2007). Because of this scale and generalization, border simplification impacts attribute measurements that are derived from the boundary data, especially for small areas and at the local scale. In particular, percentages (unitless ratios) generated from attribute measurements made for the smallest communities or county subdivisions may be under- or overstated depending upon the relative location of the smoothed border of the geopolitical unit.

While the 2001 NLCD is a substantial improvement over the 1991 AVHRR data (30-m versus 1-km resolution), it also has local-scale data and application

limitations. Initial tree canopy cover results revealed mean absolute errors (mean of the absolute difference between predicted and actual values) from 8.4 percent to 14.1 percent, with correlation coefficients between predicted and actual values ranging from 0.78 to 0.93. Impervious surface cover results revealed mean absolute errors from 4.6 percent to 7 percent, with *r*-values from 0.83 to 0.91 (Homer et al. 2004).

A more recent analysis of 127 community and 20 county geographies sampled throughout the continental United States compared NLCD tree canopy and impervious surface cover estimates with high resolution (1-m or less resolution) aerial photo-interpreted estimates. This analysis revealed that NLCD underestimates both tree canopy and impervious surface cover compared to photo-interpreted values. NLCD underestimates of tree cover vary by mapping zone, while underestimates of impervious surface cover, which are relatively minor, varies by population density (Greenfield et al. 2009). These findings are consistent with Walton (2008), who found a consistent under-prediction bias for the 2001 NLCD derived tree canopy cover values in census places (communities) of western New York.

The tree cover and impervious cover data given in this report are directly from the NLCD database. To help understand the potential underestimate in the cover values, each U.S. mapping zone was photo-interpreted using Google Earth images³. Table B provides a comparison of results from NLCD versus photo-interpreted data for mapping zones applicable to this collection of states.

Comparisons between NLCD impervious surface cover estimates and photo-interpreted values were not reported because differences were related to population density, which can vary significantly among geographic units. Despite the potential underestimates in tree canopy cover values, relative comparisons

³Nowak, D.J.; Greenfield, E.J. Tree and impervious cover in the conterminous United States: Testing of NLCD cover estimates by mapping zone. In review.

Table B.—Comparison of NLCD versus photo-interpretation (PI) derived values of percent tree canopy cover by NLCD mapping zones

Mapping zone ^a	n ^b	Percent tree canopy cover		Difference ^e	Margin of error ^f	Significant difference ^g
		NLCD ^c	PI ^d			
41	885	50.1%	53.2%	3.0%	2.6%	Yes
42	452	7.8%	10.5%	2.7%	2.0%	Yes
44	913	50.0%	60.6%	10.7%	2.1%	Yes
47	451	30.5%	41.1%	10.6%	2.9%	Yes
49	350	12.1%	15.0%	2.9%	3.1%	No
50	241	36.1%	44.2%	8.1%	4.5%	Yes
51	381	41.8%	56.4%	14.6%	3.6%	Yes
52	846	9.3%	15.5%	6.3%	1.9%	Yes
53	223	66.8%	77.2%	10.4%	4.1%	Yes
62	845	50.0%	59.7%	9.7%	2.6%	Yes

^a NLCD mapping zones.

^b Number of photo-interpreted sample points

^c Percent tree canopy value derived from NLCD data

^d Percent tree canopy derived from photo-interpreted data

^e PI value minus NLCD value

^f 95% confidence interval of PI value

^g Significant difference between NLCD and PI values if NLCD value is outside of 95% confidence interval of PI value

of tree cover among geographies in this report (e.g., planting priority index and the ratings of excellent to poor for local government tree cover) are reasonable as the under-prediction of tree cover is fairly consistent within each mapping zone. However, it is important to note that the tree canopy and impervious surface cover could be underestimated, as well as their associated ecosystem services and values. A forthcoming analysis will better assess the accuracy of the NLCD cover maps (Homer et al. 2007), but these maps and data provide comprehensive, consistent, and comparable estimates (with an inherent degree of error and uncertainty) of tree canopy and impervious surface cover to help urban and community forest management, planning and policy making. Higher resolution cover data may provide more accurate results at the local scale, but the NLCD cover maps provide a cost-effective means to consistently assess and compare the relative differences of urban cover types regionally. For more refined and locally appropriate data, local field or high resolution (1 m or less) image analyses are recommended (e.g., i-Tree www.itreetools.org; UTC – www.nrs.fs.fed.us/urban/utc).

Because of limited urban and community forest field data, data from several urban and community forests

were used to estimate the number of trees and carbon storage by trees. These coarse estimates reveal that urban and community forests contain a large number of trees and provide significant environmental benefits. Field data are needed from all states to help improve these estimates as well as to estimate other forest effects (e.g., building energy conservation and changes in stream flow and water quality). Data from long-term monitoring of urban and community forests used in conjunction with satellite-based cover maps will provide essential information to assess forest health and change, and to improve urban and community forest management.

Practical Applications for Managers

The data from this report can be used to aid urban forest management at both the state and local levels.

Data can be used to:

- Determine the extent, magnitude, and variation in the urban and community forestry resource
- Determine areas of greatest population growth, urbanization, and development (sprawl) to direct urban and community forestry to minimize negative impacts and maximize environmental benefits


- Evaluate existing tree canopy, impervious surface cover, and available planting space (available green space) to direct current and future urban and community forestry efforts such as planting programs
- Compare tree canopy cover for similar geopolitical units and set tree canopy goals
- Prioritize tree planting based on population density, tree canopy green space, and tree canopy cover per person
- Understand the pollution removal and carbon storage benefits of urban and community forests
- Promote more detailed and/or locally appropriate urban and community forest inventories, censuses, or field surveys (e.g. i-Tree - www.itreetools.org)
- Establish statewide to local standards related to urban and community forestry (e.g., establishing minimum goals of percent canopy green space or tree cover per capita and directing resources so that communities can reach the minimum standards)
- Improve urban and community forest management and cost estimation by providing an estimate of the number of trees in each geopolitical unit (i.e., urban area size (ha) * percent tree cover * 477 trees/ha, or local tree density information from local data)
- Guide policy decisions related to urban sprawl and urban and community forest management

SUMMARY

The data presented in this report provide a better understanding of urban and community forests. This information can be used to advance urban and community forest policy and management that could improve environmental quality and human health throughout the state. The following sections detail specific urban and community forestry data for the states in this regional report.

ACKNOWLEDGMENTS

This research was funded, in part, by the U.S. Forest Service's RPA Assessment Staff, State and Private Forestry's Urban and Community Forestry Program, and Northeastern Area State and Private Forestry. Thanks also goes to Chris Sorrentino for assistance with report compilation, Nana Efua Imbeah for assistance with data processing, and Mike Boarman for assistance with image processing.



ILLINOIS' URBAN AND COMMUNITY FORESTS

Statewide Summary

Urban or community land in Illinois comprises about 8.7 percent of the state land area in 2000, an increase from 7.5 percent in 1990. Statewide tree canopy cover averages 12.1 percent and tree cover in urban or community areas is about 13.0 percent, with 25.9 percent impervious surface cover and 17.5 percent of the total green space covered by tree canopy cover. Statewide, urban or community land in Illinois has an estimated 77 million trees, which store about 14.7 million metric tons of carbon (\$335.2 million), and annually remove about 484,000 metric tons of carbon (\$11.0 million) and 13,560 metric tons of air pollution (\$107.9 million) (Table IL-1).

Tables IL-2 through IL-17 are not printed in this report but are available on the CD located on the inside back cover and at <http://nrs.fs.fed.us/data/urban>.

Table IL-1.—Statewide summary of population, area, population density, tree canopy and impervious surface land cover, and urban tree benefits in urban, community, and urban or community areas.

Illinois		Statewide	Urban ^a	Community ^b	Urban or Community ^c	
Population	2000	12,419,293	10,909,520	10,749,236	n/a	
	1990	11,430,602	9,668,552	9,726,060	n/a	
	% Change (1990-2000)	8.6	12.8	10.5	n/a	
	% Total population (2000)	100.0	87.8	86.6	n/a	
Total area	km ² (2000)	149,997.6	9,337.2	10,380.4	12,796.8	
	km ² (1990)	149,997.6	8,070.7	8,903.2	11,121.3	
	% Change (1990-2000)	0.0	15.7	16.6	15.1	
Land area	km ² (2000)	143,526.6	9,199.1	10,084.7	12,452.3	
	% Land area (2000)	100.0	6.4	7.0	8.7	
	km ² (1990)	143,526.6	7,963.1	8,649.8	10,824.6	
	% Land area (1990)	100.0	5.5	6.0	7.5	
	% Change (1990-2000)	0.0	15.5	16.6	15.0	
Population density (people/land area km ²)	2000	86.5	1,185.9	1,065.9	n/a	
	1990	79.6	1,214.2	1,124.4	n/a	
	% Change (1990-2000)	8.6	-2.3	-5.2	n/a	
Tree canopy cover (2000)	km ²	17,361.5	1,180.2	1,168.4	1,613.8	
	% Land area	12.1	12.8	11.6	13.0	
	Per capita (m ² /person)	1,397.9	108.2	108.7	n/a	
	% Canopy green space ^d	12.5	18.4	16.3	17.5	
Total green space (2000) ^e	km ²	138,831.0	6,399.9	7,147.7	9,224.3	
	% Land area	96.7	69.6	70.9	74.1	
Available green space (2000) ^f	km ²	121,472.0	5,220.9	5,980.4	7,611.8	
	% Land area	84.6	56.8	59.3	61.1	
Impervious surface cover (2000)	km ²	4,695.2	2,799.2	2,937.0	3,228.0	
	% Land area	3.3	30.4	29.1	25.9	
	Per capita (m ² /person)	378.1	256.6	273.2	n/a	
Urban tree benefits (2000)	Estimated number of trees	n/a	56,300,000	55,700,000	77,000,000	
	Carbon					
	Carbon stored (metric tons)	n/a	10,700,000	10,600,000	14,700,000	
	Carbon stored (\$)	n/a	\$244,000,000	\$241,700,000	\$335,200,000	
	Carbon sequestered (metric tons/year)	n/a	354,000	351,000	484,000	
	Carbon sequestered (\$/year)	n/a	\$8,071,000	\$8,003,000	\$11,035,000	
	Pollution					
	CO removed (metric tons/year)	n/a	171	170	234	
	CO removed (\$/year)	n/a	\$241,100	\$238,700	\$329,700	
	NO ₂ removed (metric tons/year)	n/a	1,936	1,916	2,647	
	NO ₂ removed (\$/year)	n/a	\$19,173,400	\$18,981,600	\$26,217,300	
	O ₃ removed (metric tons/year)	n/a	3,466	3,431	4,739	
	O ₃ removed (\$/year)	n/a	\$34,334,000	\$33,990,000	\$46,947,000	
	SO ₂ removed (metric tons/year)	n/a	851	843	1,164	
	SO ₂ removed (\$/year)	n/a	\$2,064,800	\$2,044,100	\$2,823,300	
PM ₁₀ removed (metric tons/year)	n/a	3,490	3,455	4,772		
PM ₁₀ removed (\$/year)	n/a	\$23,081,200	\$22,850,200	\$31,560,300		
Total pollution removal (metric tons/year)	n/a	9,910	9,810	13,560		
Total pollution removal (\$/year)	n/a	\$78,900,000	\$78,100,000	\$107,900,000		

^a Urban land is based on population density and was delimited using the United States Census definitions of urbanized areas and urban clusters. ^b Community land is based on jurisdictional or political boundaries of communities based on United States Census definitions of incorporated or census designated places. ^c Urban or communities is land that is urban, community, or both. Communities may include all, some, or no urban land within their boundaries. ^d Canopy green space is the tree canopy cover divided by total green space. ^e Total green space (TGS) is total area – impervious surface cover – water. ^f Available green space (AGS) is total green space – tree canopy cover (if the calculated value is less than 0, then value set at 0).




Figure IL-1.—Urban or community land in 2000; urban area relative to community boundaries.




Figure IL-2.—2000 population within county subdivision boundaries.

Human Population Characteristics and Trends

The population in Illinois increased 8.6 percent, from 11,430,602 in 1990 to 12,419,293 in 2000 (Table IL-1). In Illinois, 87.8 percent of the State's population is in urban areas (Fig. IL-1), and 86.6 percent of the population is within communities (Fig. IL-2).

Urban and Community Land

Urban land comprises 6.4 percent of the land area of Illinois, while lands within communities make up 7.0 percent of the State (Fig. IL-1). Between 1990 and 2000, urban area increased 15.5 percent, while community land increased from 6.0 to 7.0 percent (Table IL-1). Urban area in Illinois is projected to increase to 14.6 percent by 2050, based on average urban growth pattern of the 1990s (Nowak and Walton 2005). Both urban land (attaining minimum population density) and community land (political boundaries) increased from 1990 to 2000. The percentages are calculated using the total (water and land) area of the geopolitical units derived from U.S. Census cartographic boundary data. Percent urban land varied across the State (Fig. IL-3; Tables IL-2 through 4).




Figure IL-3.—Percent of county subdivision area classified as urban land in 2000.




Figure IL-4.—Percentage tree canopy cover.




Figure IL-5.—Percentage tree canopy cover within county subdivisions.

Tree Canopy Cover Characteristics

Tree canopy cover in Illinois averages 12.1 percent (Fig. IL-4), with 96.7 percent total green space, 12.5 percent canopy green space, and 1,397.9 m² of canopy cover per capita. Average tree cover in urban areas in Illinois was 12.8 percent, with 69.6 percent total green space, 18.4 percent canopy green space, and 108.2 m² of canopy cover per capita. Within community lands in Illinois, average tree cover was 11.6 percent, with 70.9 percent total green space, 16.3 percent canopy green space, and 108.7 m² of canopy cover per capita (Table IL-1). Tree canopy cover, canopy green space, and tree cover per capita varied among communities, county subdivisions, and counties (Fig. IL-5 through 6; Tables IL-5 through 7).




Figure IL-6.—Percentage tree canopy green space in county subdivisions.




Figure IL-7.—Percentage impervious surface cover.




Figure IL-8.—Percentage impervious surface cover within county subdivisions.

Impervious Surface Cover Characteristics

Average impervious surface cover in Illinois is 3.3 percent of the land area (Fig. IL-7), with 378.1 m² of impervious surface cover per capita. Average impervious surface cover in urban areas was 30.4 percent, with 256.6 m² of impervious surface cover per capita. Within community lands in Illinois, average impervious surface cover was 29.1 percent with 273.2 m² of impervious surface cover per capita (Table IL-1). Impervious surface cover varied across the state (Fig. IL-8; Tables IL-5 through 7).




Figure IL-9.—Classified land cover.




Figure IL-10.—Relative comparisons of tree cover for county subdivisions.

Classified Land-cover Characteristics

Illinois' land cover is dominated by agricultural land (Fig. IL-9). The characteristics as a percent of the total land area in Illinois are (Tables IL-8 through 10):

- Agricultural – 71.5 percent
- Forested – 15.6 percent
- Developed – 11.5 percent
- Grassland – 0.8 percent
- Wetland – 0.2 percent
- Barren – 0.1 percent
- Scrub/Shrub – Less than 0.1 percent

Relative Comparisons of Tree Cover

Out of the 1,313 Illinois communities, eight received a rating of excellent and 1,121 received a rating of poor (Table IL-12). Of the 1,708 county subdivisions, eight had a rating of excellent and 1,332 were rated poor (Fig. IL-10, Table IL-13); and out of 102 counties, five were given a rating of excellent and 68 were given a rating of poor (Table IL-14). Variability of assessment scores is a product of the difference in land cover distributions and the percentage of canopy cover within the population density classes and mapping zones (Fig. IL-10; Tables IL-11 through 14).




Figure IL-11.—Planting priority index for county subdivisions. The higher the index value, the greater priority for planting.


Priority Areas for Tree Planting

Priority areas for planting tend to be highest in more urbanized areas due to higher population density (Fig. IL-11; Tables IL-15 through 17). These index values can also be produced using high resolution cover data to determine local planting priority areas (e.g., neighborhoods).

Urban Tree Benefits

The following forest attributes are estimated for the urban or community land in Illinois (Table IL-1). These are rough estimates of values. More localized data are needed for more precise estimates, but these values reveal first-order approximations.

- 77 million trees
- 14.7 million metric tons of C stored (\$335.2 million value)
- 484,000 metric tons/year of C sequestered (\$11.0 million value)
- 13,560 metric tons/year total pollution removal (\$107.9 million value)
 - 234 metric tons/year of CO removed (\$329,700 value)
 - 2,647 metric tons/year NO₂ removed (\$26.2 million value)
 - 4,739 metric tons/year of O₃ removed (\$46.9 million value)
 - 1,164 metric tons/year of SO₂ removed (\$2.8 million value)
 - 4,772 metric tons/year of PM₁₀ removed (\$31.6 million value)




Summary

The data presented in this report provide a better understanding of Illinois' urban and community forests. This information can be used to advance urban and community forest policy and management that could improve environmental quality and human health throughout the State.

These data establish a baseline to assess future change and can be used to understand:

- Extent of the urban and community forest resource
- Variations in the resource across the State
- Magnitude and value of the urban and community forest resource
- Urban growth in Illinois
- Implications of policy decisions related to urban sprawl and urban and community forest management



INDIANA'S URBAN AND COMMUNITY FORESTS

Statewide Summary

Urban or community land in Indiana comprises about 8.1 percent of the state land area in 2000, an increase from 6.9 percent in 1990. Statewide tree canopy cover averages 18.2 percent and tree cover in urban or community areas is about 14.5 percent, with 19.1 percent impervious surface cover and 17.9 percent of the total green space covered by tree canopy cover. Statewide, urban or community land in Indiana has an estimated 52 million trees, which store about 9.9 million metric tons of carbon (\$225.7 million), and annually remove about 327,000 metric tons of carbon (\$7.5 million) and 8,620 metric tons of air pollution (\$70.9 million) (Table IN-1).

Tables IN-2 through IN-17 are not printed in this report but are available on the CD located on the inside back cover and at <http://nrs.fs.fed.us/data/urban>.

Table IN-1.—Statewide summary of population, area, population density, tree canopy and impervious surface land cover, and urban tree benefits in urban, community, and urban or community areas.

Indiana		Statewide	Urban ^a	Community ^b	Urban or Community ^c	
Population	2000	6,080,485	4,304,011	3,998,963	n/a	
	1990	5,544,159	3,598,099	2,890,061	n/a	
	% Change (1990-2000)	9.7	19.6	38.4	n/a	
	% Total population (2000)	100.0	70.8	65.8	n/a	
Total area	km ² (2000)	94,321.5	5,762.1	5,847.8	7,690.7	
	km ² (1990)	94,321.5	4,814.2	4,985.3	6,523.0	
	% Change (1990-2000)	0.0	19.7	17.3	17.9	
Land area	km ² (2000)	92,654.7	5,668.1	5,723.9	7,535.0	
	% Land area (2000)	100.0	6.1	6.2	8.1	
	km ² (1990)	92,654.7	4,741.9	4,881.9	6,395.6	
	% Land area (1990)	100.0	5.1	5.3	6.9	
	% Change (1990-2000)	0.0	19.5	17.2	17.8	
Population density (people/land area km ²)	2000	65.6	759.3	698.6	n/a	
	1990	59.8	758.8	592.0	n/a	
	% Change (1990-2000)	9.7	0.1	18.0	n/a	
Tree canopy cover (2000)	km ²	16,856.9	822.0	768.7	1,090.0	
	% Land area	18.2	14.5	13.4	14.5	
	Per capita (m ² /person)	2,772.3	191.0	192.2	n/a	
	% Canopy green space ^d	18.6	18.6	17.3	17.9	
Total green space (2000) ^e	km ²	90,565.2	4,418.8	4,443.3	6,095.3	
	% Land area	97.7	78.0	77.6	80.9	
Available green space (2000) ^f	km ²	73,708.9	3,597.3	3,675.0	5,005.8	
	% Land area	79.6	63.5	64.2	66.4	
Impervious surface cover (2000)	km ²	2,089.5	1,249.4	1,280.6	1,439.8	
	% Land area	2.3	22.0	22.4	19.1	
	Per capita (m ² /person)	343.6	290.3	320.2	n/a	
Urban tree benefits (2000)	Estimated number of trees	n/a	39,200,000	36,700,000	52,000,000	
	Carbon					
	Carbon stored (metric tons)	n/a	7,500,000	7,000,000	9,900,000	
	Carbon stored (\$)	n/a	\$171,000,000	\$159,600,000	\$225,700,000	
	Carbon sequestered (metric tons/year)	n/a	247,000	231,000	327,000	
	Carbon sequestered (\$/year)	n/a	\$5,632,000	\$5,267,000	\$7,456,000	
	Pollution					
	CO removed (metric tons/year)	n/a	108	101	143	
	CO removed (\$/year)	n/a	\$151,600	\$141,800	\$201,100	
	NO ₂ removed (metric tons/year)	n/a	885	828	1,174	
	NO ₂ removed (\$/year)	n/a	\$8,769,700	\$8,201,600	\$11,629,100	
	O ₃ removed (metric tons/year)	n/a	3,048	2,851	4,042	
	O ₃ removed (\$/year)	n/a	\$30,193,000	\$28,237,000	\$40,038,000	
	SO ₂ removed (metric tons/year)	n/a	461	432	612	
SO ₂ removed (\$/year)	n/a	\$1,118,900	\$1,046,400	\$1,483,700		
PM ₁₀ removed (metric tons/year)	n/a	2,002	1,872	2,654		
PM ₁₀ removed (\$/year)	n/a	\$13,239,400	\$12,381,700	\$17,556,100		
Total pollution removal (metric tons/year)	n/a	6,500	6,080	8,620		
Total pollution removal (\$/year)	n/a	\$53,500,000	\$50,000,000	\$70,900,000		

^a Urban land is based on population density and was delimited using the United States Census definitions of urbanized areas and urban clusters. ^b Community land is based on jurisdictional or political boundaries of communities based on United States Census definitions of incorporated or census designated places. ^c Urban or communities is land that is urban, community, or both. Communities may include all, some, or no urban land within their boundaries. ^d Canopy green space is the tree canopy cover divided by total green space. ^e Total green space (TGS) is total area – impervious surface cover – water. ^f Available green space (AGS) is total green space – tree canopy cover (if the calculated value is less than 0, then value set at 0).




Figure IN-1.—Urban or community land in 2000; urban area relative to community boundaries.




Figure IN-2.—2000 population within county subdivision boundaries.

Human Population Characteristics and Trends

The population in Indiana increased 9.7 percent, from 5,544,159 in 1990 to 6,080,485 in 2000 (Table IN-1). In Indiana, 70.8 percent of the State’s population is in urban areas (Fig. IN-1), and 65.8 percent of the population is within communities (Fig. IN-2).

Urban and Community Land

Urban land comprises 6.1 percent of the land area of Indiana, while lands within communities make up 6.2 percent of the State (Fig. IN-1). Between 1990 and 2000, urban area increased 19.5 percent, while community land increased from 5.3 to 6.2 percent (Table IN-1). Urban area in Indiana is projected to increase to 16.7 percent by 2050, based on average urban growth pattern of the 1990s (Nowak and Walton 2005). Both urban land (attaining minimum population density) and community land (political boundaries) increased from 1990 to 2000. The percentages are calculated using the total (water and land) area of the geopolitical units derived from U.S. Census cartographic boundary data. Percent urban land varied across the State (Fig. IN-3; Tables IN-2 through 4).




Figure IN-3.—Percent of county subdivision area classified as urban land in 2000.




Figure IN-4.—Percentage tree canopy cover.




Figure IN-5.—Percentage tree canopy cover within county subdivisions.

Tree Canopy Cover Characteristics

Tree canopy cover in Indiana averages 18.2 percent (Fig. IN-4), with 97.7 percent total green space, 18.6 percent canopy green space, and 2,772.3 m² of canopy cover per capita. Average tree cover in urban areas in Indiana was 14.5 percent, with 78.0 percent total green space, 18.6 percent canopy green space, and 191.0 m² of canopy cover per capita. Within community lands in Indiana, average tree cover was 13.4 percent, with 77.6 percent total green space, 17.3 percent canopy green space, and 192.2 m² of canopy cover per capita (Table IN-1). Tree canopy cover, canopy green space, and tree cover per capita varied among communities, county subdivisions, and counties (Fig. IN-5 through 6; Tables IN-5 through 7).




Figure IN-6.—Percentage tree canopy green space in county subdivisions.




Figure IN-7.—Percentage impervious surface cover.




Figure IN-8.—Percentage impervious surface cover within county subdivisions.

Impervious Surface Cover Characteristics

Average impervious surface cover in Indiana is 2.3 percent of the land area (Fig. IN-7), with 343.6 m² of impervious surface cover per capita. Average impervious surface cover in urban areas was 22.0 percent, with 290.3 m² of impervious surface cover per capita. Within community lands in Indiana, average impervious surface cover was 22.4 percent with 320.2 m² of impervious surface cover per capita (Table IN-1). Impervious surface cover varied across the state (Fig. IN-8; Tables IN-5 through 7).




Figure IN-9.—Classified land cover.




Figure IN-10.—Relative comparisons of tree cover for county subdivisions.

Classified Land-cover Characteristics

Indiana’s land cover is dominated by agricultural land (Fig. IN-9). The characteristics as a percent of the total land area in Indiana are (Tables IN-8 through 10):

- Agricultural – 64.3 percent
- Forested – 23.2 percent
- Developed – 10.4 percent
- Grassland – 1.6 percent
- Scrub/Shrub – 0.3 percent
- Wetland – 0.2 percent
- Barren – 0.1 percent

Relative Comparisons of Tree Cover

Out of the 601 Indiana communities, 12 received a rating of excellent and 465 received a rating of poor (Table IN-12). Of the 1,009 county subdivisions, 10 had a rating of excellent and 637 were rated poor (Fig. IN-10, Table IN-13); and out of 92 counties, four were given a rating of excellent and 45 were given a rating of poor (Table IN-14). Variability of assessment scores is a product of the difference in land cover distributions and the percentage of canopy cover within the population density classes and mapping zones (Fig. IN-10; Tables IN-11 through 14).




Figure IN-11.—Planting priority index for county subdivisions. The higher the index value, the greater priority for planting.


Priority Areas for Tree Planting

Priority areas for planting tend to be highest in more urbanized areas due to higher population density (Fig. IN-11; Tables IN-15 through 17). These index values can also be produced using high resolution cover data to determine local planting priority areas (e.g., neighborhoods).

Urban Tree Benefits

The following forest attributes are estimated for the urban or community land in Indiana (Table IN-1). These are rough estimates of values. More localized data are needed for more precise estimates, but these values reveal first-order approximations.

- 52 million trees
- 9.9 million metric tons of C stored (\$225.7 million value)
- 327,000 metric tons/year of C sequestered (\$7.5 million value)
- 8,620 metric tons/year total pollution removal (\$70.9 million value)
 - 143 metric tons/year of CO removed (\$201,100 value)
 - 1,174 metric tons/year NO₂ removed (\$11.6 million value)
 - 4,042 metric tons/year of O₃ removed (\$40.0 million value)
 - 612 metric tons/year of SO₂ removed (\$1.5 million value)
 - 2,654 metric tons/year of PM₁₀ removed (\$17.6 million value).




Summary

The data presented in this report provide a better understanding of Indiana's urban and community forests. This information can be used to advance urban and community forest policy and management that could improve environmental quality and human health throughout the State.

These data establish a baseline to assess future change and can be used to understand:

- Extent of the urban and community forest resource
- Variations in the resource across the State
- Magnitude and value of the urban and community forest resource
- Urban growth in Indiana
- Implications of policy decisions related to urban sprawl and urban and community forest management



MICHIGAN'S URBAN AND COMMUNITY FORESTS

Statewide Summary

Urban or community land in Michigan comprises about 7.3 percent of the state land area in 2000, an increase from 6.5 percent in 1990. Statewide tree canopy cover averages 42.9 percent and tree cover in urban or community areas is about 21.0 percent, with 24.2 percent impervious surface cover and 27.8 percent of the total green space covered by tree canopy cover. Statewide, urban or community land in Michigan has an estimated 107.8 million trees, which store about 20.6 million metric tons of carbon (\$469.7 million), and annually remove about 678,000 metric tons of carbon (\$15.5 million) and 14,820 metric tons of air pollution (\$121.7 million) (Table MI-1).

Tables MI-2 through MI-17 are not printed in this report but are available on the CD located on the inside back cover and at <http://nrs.fs.fed.us/data/urban>.

Table MI-1.—Statewide summary of population, area, population density, tree canopy and impervious surface land cover, and urban tree benefits in urban, community, and urban or community areas.

Michigan		Statewide	Urban ^a	Community ^b	Urban or Community ^c	
Population	2000	9,938,444	7,419,457	6,384,273	n/a	
	1990	9,295,297	6,555,842	6,073,842	n/a	
	% Change (1990-2000)	6.9	13.2	5.1	n/a	
	% Total population (2000)	100.0	74.7	64.2	n/a	
Total area	km ² (2000)	250,493.6	8,879.5	7,871.4	11,283.6	
	km ² (1990)	250,493.6	7,576.6	7,417.3	9,933.7	
	% Change (1990-2000)	0.0	17.2	6.1	13.6	
Land area	km ² (2000)	146,909.8	8,604.2	7,449.4	10,740.5	
	% Land area (2000)	100.0	5.9	5.1	7.3	
	km ² (1990)	146,909.8	7,403.7	7,093.1	9,537.1	
	% Land area (1990)	100.0	5.0	4.8	6.5	
	% Change (1990-2000)	0.0	16.2	5.0	12.6	
Population density (people/land area km ²)	2000	67.6	862.3	857.0	n/a	
	1990	63.3	885.5	856.3	n/a	
	% Change (1990-2000)	6.9	-2.6	0.1	n/a	
Tree canopy cover (2000)	km ²	63,024.9	1,629.0	1,456.2	2,259.8	
	% Land area	42.9	18.9	19.5	21.0	
	Per capita (m ² /person)	6,341.5	219.6	228.1	n/a	
	% Canopy green space ^d	44.1	26.2	27.3	27.8	
Total green space (2000) ^e	km ²	142,933.0	6,223.4	5,332.5	8,137.1	
	% Land area	97.3	72.3	71.6	75.8	
Available green space (2000) ^f	km ²	79,910.2	4,595.9	3,877.7	5,878.9	
	% Land area	54.4	53.4	52.1	54.7	
Impervious surface cover (2000)	km ²	3,977.0	2,380.8	2,116.9	2,603.5	
	% Land area	2.7	27.7	28.4	24.2	
	Per capita (m ² /person)	400.2	320.9	331.6	n/a	
Urban tree benefits (2000)	Estimated number of trees	n/a	77,700,000	69,400,000	107,800,000	
	Carbon					
	Carbon stored (metric tons)	n/a	14,800,000	13,300,000	20,600,000	
	Carbon stored (\$)	n/a	\$337,400,000	\$303,200,000	\$469,700,000	
	Carbon sequestered (metric tons/year)	n/a	489,000	437,000	678,000	
	Carbon sequestered (\$/year)	n/a	\$11,149,000	\$9,964,000	\$15,458,000	
	Pollution					
	CO removed (metric tons/year)	n/a	113	101	157	
	CO removed (\$/year)	n/a	\$159,700	\$142,700	\$221,500	
	NO ₂ removed (metric tons/year)	n/a	1,753	1,567	2,432	
	NO ₂ removed (\$/year)	n/a	\$17,369,400	\$15,527,500	\$24,095,600	
	O ₃ removed (metric tons/year)	n/a	4,588	4,101	6,364	
	O ₃ removed (\$/year)	n/a	\$45,445,000	\$40,626,000	\$63,044,000	
	SO ₂ removed (metric tons/year)	n/a	764	683	1,060	
	SO ₂ removed (\$/year)	n/a	\$1,852,300	\$1,655,900	\$2,569,600	
PM ₁₀ removed (metric tons/year)	n/a	3,464	3,097	4,806		
PM ₁₀ removed (\$/year)	n/a	\$22,912,300	\$20,482,700	\$31,785,000		
Total pollution removal (metric tons/year)	n/a	10,680	9,550	14,820		
Total pollution removal (\$/year)	n/a	\$87,700,000	\$78,400,000	\$121,700,000		

^a Urban land is based on population density and was delimited using the United States Census definitions of urbanized areas and urban clusters. ^b Community land is based on jurisdictional or political boundaries of communities based on United States Census definitions of incorporated or census designated places. ^c Urban or communities is land that is urban, community, or both. Communities may include all, some, or no urban land within their boundaries. ^d Canopy green space is the tree canopy cover divided by total green space. ^e Total green space (TGS) is total area – impervious surface cover – water. ^f Available green space (AGS) is total green space – tree canopy cover (if the calculated value is less than 0, then value set at 0).




Figure MI-1.—Urban or community land in 2000; urban area relative to community boundaries.




Figure MI-2.—2000 population within county subdivision boundaries.

Human Population Characteristics and Trends

The population in Michigan increased 6.9 percent, from 9,295,297 in 1990 to 9,938,444 in 2000 (Table MI-1). In Michigan, 74.7 percent of the State’s population is in urban areas (Fig. MI-1), and 64.2 percent of the population is within communities (Fig. MI-2).

Urban and Community Land

Urban land comprises 5.9 percent of the land area of Michigan, while lands within communities make up 5.1 percent of the State (Fig. MI-1). Between 1990 and 2000, urban area increased 16.2 percent, while community land increased from 4.8 to 5.1 percent (Table MI-1). Urban area in Michigan is projected to increase to 13.7 percent by 2050, based on average urban growth pattern of the 1990s (Nowak and Walton 2005). Both urban land (attaining minimum population density) and community land (political boundaries) increased from 1990 to 2000. The percentages are calculated using the total (water and land) area of the geopolitical units derived from U.S. Census cartographic boundary data. Percent urban land varied across the State (Fig. MI-3; Tables MI-2 through 4).




Figure MI-3.—Percent of county subdivision area classified as urban land in 2000.




Figure MI-4.—Percentage tree canopy cover.




Figure MI-5.—Percentage tree canopy cover within county subdivisions.

Tree Canopy Cover Characteristics

Tree canopy cover in Michigan averages 42.9 percent (Fig. MI-4), with 97.3 percent total green space, 44.1 percent canopy green space, and 6,341.5 m² of canopy cover per capita. Average tree cover in urban areas in Michigan was 18.9 percent, with 72.3 percent total green space, 26.2 percent canopy green space, and 219.6 m² of canopy cover per capita. Within community lands in Michigan, average tree cover was 19.5 percent, with 71.6 percent total green space, 27.3 percent canopy green space, and 228.1 m² of canopy cover per capita (Table MI-1). Tree canopy cover, canopy green space, and tree cover per capita varied among communities, county subdivisions, and counties (Fig. MI-5 through 6; Tables MI-5 through 7).




Figure MI-6.—Percentage tree canopy green space in county subdivisions.




Figure MI-7.—Percentage impervious surface cover.




Figure MI-8.—Percentage impervious surface cover within county subdivisions.

Impervious Surface Cover Characteristics

Average impervious surface cover in Michigan is 2.7 percent of the land area (Fig. MI-7), with 400.2 m² of impervious surface cover per capita. Average impervious surface cover in urban areas was 27.7 percent, with 320.9 m² of impervious surface cover per capita. Within community lands in Michigan, average impervious surface cover was 28.4 percent with 331.6 m² of impervious surface cover per capita (Table MI-1). Impervious surface cover varied across the state (Fig. MI-8; Tables MI-5 through 7).




Figure MI-9.—Classified land cover.




Figure MI-10.—Relative comparisons of tree cover for county subdivisions.

Classified Land-cover Characteristics

Michigan’s land cover is dominated by agricultural land (Fig. MI-9). The characteristics as a percent of the total land area in Michigan are (Tables MI-8 through 10):

- Agricultural – 44.4 percent
- Forested – 36.5 percent
- Developed – 10.8 percent
- Grassland – 5.0 percent
- Wetland – 1.7 percent
- Scrub/Shrub – 1.0 percent
- Barren – 0.5 percent

Relative Comparisons of Tree Cover

Out of the 630 Michigan communities, 15 received a rating of excellent and 289 received a rating of poor (Table MI-12). Of the 1,529 county subdivisions, 38 had a rating of excellent and 543 were rated poor (Fig. MI-10, Table MI-13); and out of 83 counties, 12 were given a rating of excellent and 18 were given a rating of poor (Table MI-14). Variability of assessment scores is a product of the difference in land cover distributions and the percentage of canopy cover within the population density classes and mapping zones (Fig. MI-10; Tables MI-11 through 14).




Figure MI-11.—Planting priority index for county subdivisions. The higher the index value, the greater priority for planting.


Priority Areas for Tree Planting

Priority areas for planting tend to be highest in more urbanized areas due to higher population density (Fig. MI-11; Tables MI-15 through 17). These index values can also be produced using high resolution cover data to determine local planting priority areas (e.g., neighborhoods).

Urban Tree Benefits

The following forest attributes are estimated for the urban or community land in Michigan (Table MI-1). These are rough estimates of values. More localized data are needed for more precise estimates, but these values reveal first-order approximations.

- 107.8 million trees
- 20.6 million metric tons of C stored (\$469.7 million value)
- 678,000 metric tons/year of C sequestered (\$15.5 million value)
- 14,820 metric tons/year total pollution removal (\$121.7 million value)
 - 157 metric tons/year of CO removed (\$221,500 value)
 - 2,432 metric tons/year NO₂ removed (\$24.1 million value)
 - 6,364 metric tons/year of O₃ removed (\$63.0 million value)
 - 1,060 metric tons/year of SO₂ removed (\$2.6 million value)
 - 4,806 metric tons/year of PM₁₀ removed (\$31.8 million value)




Summary

The data presented in this report provide a better understanding of Michigan's urban and community forests. This information can be used to advance urban and community forest policy and management that could improve environmental quality and human health throughout the State.

These data establish a baseline to assess future change and can be used to understand:

- Extent of the urban and community forest resource
- Variations in the resource across the State
- Magnitude and value of the urban and community forest resource
- Urban growth in Michigan
- Implications of policy decisions related to urban sprawl and urban and community forest management



OHIO'S URBAN AND COMMUNITY FORESTS

Statewide Summary

Urban or community land in Ohio comprises about 12.6 percent of the state land area in 2000, an increase from 11.4 percent in 1990. Statewide tree canopy cover averages 28.7 percent and tree cover in urban or community areas is about 21.0 percent, with 19.6 percent impervious surface cover and 26.2 percent of the total green space covered by tree canopy cover. Statewide, urban or community land in Ohio has an estimated 133.5 million trees, which store about 25.5 million metric tons of carbon (\$581.4 million), and annually remove about 840,000 metric tons of carbon (\$19.2 million) and 21,930 metric tons of air pollution (\$178.1 million) (Table OH-1).

Tables OH-2 through OH-17 are not printed in this report but are available on the CD located on the inside back cover and at <http://nrs.fs.fed.us/data/urban>.

Table OH-1.—Statewide summary of population, area, population density, tree canopy and impervious surface land cover, and urban tree benefits in urban, community, and urban or community areas.

Ohio		Statewide	Urban ^a	Community ^b	Urban or Community ^c	
Population	2000	11,353,140	8,782,329	8,012,241	n/a	
	1990	10,847,115	8,039,409	7,699,111	n/a	
	% Change (1990-2000)	4.7	9.2	4.1	n/a	
	% Total population (2000)	100.0	77.4	70.6	n/a	
Total area	km ² (2000)	116,096.0	10,402.5	10,417.4	13,570.7	
	km ² (1990)	116,096.0	9,339.0	9,346.4	12,221.8	
	% Change (1990-2000)	0.0	11.4	11.5	11.0	
Land area	km ² (2000)	105,716.9	10,272.8	10,214.4	13,318.3	
	% Land area (2000)	100.0	9.7	9.7	12.6	
	km ² (1990)	105,716.9	9,239.8	9,168.8	12,005.5	
	% Land area (1990)	100.0	8.7	8.7	11.4	
	% Change (1990-2000)	0.0	11.2	11.4	10.9	
Population density (people/land area km ²)	2000	107.4	854.9	784.4	n/a	
	1990	102.6	870.1	839.7	n/a	
	% Change (1990-2000)	4.7	-1.7	-6.6	n/a	
Tree canopy cover (2000)	km ²	30,380.8	2,030.5	2,047.4	2,799.9	
	% Land area	28.7	19.8	20.0	21.0	
	Per capita (m ² /person)	2,676.0	231.2	255.5	n/a	
	% Canopy green space ^d	29.7	25.5	25.7	26.2	
Total green space (2000) ^e	km ²	102,189.0	7,959.3	7,963.3	10,705.2	
	% Land area	96.7	77.5	78.0	80.4	
Available green space (2000) ^f	km ²	71,811.7	5,931.7	5,918.6	7,908.6	
	% Land area	67.9	57.7	57.9	59.4	
Impervious surface cover (2000)	km ²	3,528.4	2,313.5	2,251.1	2,613.1	
	% Land area	3.3	22.5	22.0	19.6	
	Per capita (m ² /person)	310.8	263.4	281.0	n/a	
Urban tree benefits (2000)	Estimated number of trees	n/a	96,800,000	97,600,000	133,500,000	
	Carbon					
	Carbon stored (metric tons)	n/a	18,500,000	18,600,000	25,500,000	
	Carbon stored (\$)	n/a	\$421,800,000	\$424,100,000	\$581,400,000	
	Carbon sequestered (metric tons/year)	n/a	609,000	614,000	840,000	
	Carbon sequestered (\$/year)	n/a	\$13,885,000	\$13,999,000	\$19,152,000	
	Pollution					
	CO removed (metric tons/year)	n/a	226	228	311	
	CO removed (\$/year)	n/a	\$317,600	\$320,300	\$438,000	
	NO ₂ removed (metric tons/year)	n/a	2,779	2,802	3,832	
	NO ₂ removed (\$/year)	n/a	\$27,532,000	\$27,761,000	\$37,963,800	
	O ₃ removed (metric tons/year)	n/a	6,641	6,696	9,157	
	O ₃ removed (\$/year)	n/a	\$65,783,000	\$66,330,000	\$90,708,000	
	SO ₂ removed (metric tons/year)	n/a	1,402	1,414	1,934	
	SO ₂ removed (\$/year)	n/a	\$3,400,400	\$3,428,700	\$4,688,800	
PM ₁₀ removed (metric tons/year)	n/a	4,854	4,894	6,693		
PM ₁₀ removed (\$/year)	n/a	\$32,105,000	\$32,372,000	\$44,269,500		
Total pollution removal (metric tons/year)	n/a	15,900	16,030	21,930		
Total pollution removal (\$/year)	n/a	\$129,100,000	\$130,200,000	\$178,100,000		

^a Urban land is based on population density and was delimited using the United States Census definitions of urbanized areas and urban clusters. ^b Community land is based on jurisdictional or political boundaries of communities based on United States Census definitions of incorporated or census designated places. ^c Urban or communities is land that is urban, community, or both. Communities may include all, some, or no urban land within their boundaries. ^d Canopy green space is the tree canopy cover divided by total green space. ^e Total green space (TGS) is total area – impervious surface cover – water. ^f Available green space (AGS) is total green space – tree canopy cover (if the calculated value is less than 0, then value set at 0).




Figure OH-1.—Urban or community land in 2000; urban area relative to community boundaries.




Figure OH-2.—2000 population within county subdivision boundaries.

Human Population Characteristics and Trends

The population in Ohio increased 4.7 percent, from 10,847,115 in 1990 to 11,353,140 in 2000 (Table OH-1). In Ohio, 77.4 percent of the State’s population is in urban areas (Fig. OH-1), and 70.6 percent of the population is within communities (Fig. OH-2).

Urban and Community Land

Urban land comprises 9.7 percent of the land area of Ohio, while lands within communities also make up 9.7 percent of the State (Fig. OH-1). Between 1990 and 2000, urban area increased 11.2 percent, while community land increased from 8.7 to 9.7 percent (Table OH-1). Urban area in Ohio is projected to increase to 22.9 percent by 2050, based on average urban growth pattern of the 1990s (Nowak and Walton 2005). Both urban land (attaining minimum population density) and community land (political boundaries) increased from 1990 to 2000. The percentages are calculated using the total (water and land) area of the geopolitical units derived from U.S. Census cartographic boundary data. Percent urban land varied across the State (Fig. OH-3; Tables OH-2 through 4).




Figure OH-3.—Percent of county subdivision area classified as urban land in 2000.




Figure OH-4.—Percentage tree canopy cover.




Figure OH-5.—Percentage tree canopy cover within county subdivisions.

Tree Canopy Cover Characteristics

Tree canopy cover in Ohio averages 28.7 percent (Fig. OH-4), with 96.7 percent total green space, 29.7 percent canopy green space, and 2,676.0 m² of canopy cover per capita. Average tree cover in urban areas in Ohio was 19.8 percent, with 77.5 percent total green space, 25.5 percent canopy green space, and 231.2 m² of canopy cover per capita. Within community lands in Ohio, average tree cover was 20.0 percent, with 78.0 percent total green space, 25.7 percent canopy green space, and 255.5 m² of canopy cover per capita (Table OH-1). Tree canopy cover, canopy green space, and tree cover per capita varied among communities, county subdivisions, and counties (Fig. OH-5 through 6; Tables OH-5 through 7).




Figure OH-6.—Percentage tree canopy green space in county subdivisions.




Figure OH-7.—Percentage impervious surface cover.




Figure OH-8.—Percentage impervious surface cover within county subdivisions.

Impervious Surface Cover Characteristics

Average impervious surface cover in Ohio is 3.3 percent of the land area (Fig. OH-7), with 310.8 m² of impervious surface cover per capita. Average impervious surface cover in urban areas was 22.5 percent, with 263.4 m² of impervious surface cover per capita. Within community lands in Ohio, average impervious surface cover was 22.0 percent with 281.0 m² of impervious surface cover per capita (Table OH-1). Impervious surface cover varied across the state (Fig. OH-8; Tables OH-5 through 7).




Figure OH-9.—Classified land cover.




Figure OH-10.—Relative comparisons of tree cover for county subdivisions.

Classified Land-cover Characteristics

Ohio's land cover is dominated by agricultural land (Fig. OH-9). The characteristics as a percent of the total land area in Ohio are (Tables OH-8 through 10):

- Agricultural – 51.7 percent
- Forested – 31.7 percent
- Developed – 14.3 percent
- Grassland – 1.6 percent
- Scrub/Shrub – 0.4 percent
- Wetland – 0.2 percent
- Barren – 0.1 percent

Relative Comparisons of Tree Cover

Out of the 1,054 Ohio communities, 10 received a rating of excellent and 629 received a rating of poor (Table OH-12). Of the 1,578 county subdivisions, 30 had a rating of excellent and 776 were rated poor (Fig. OH-10, Table OH-13); and out of 88 counties, three were given a rating of excellent and 48 were given a rating of poor (Table OH-14). Variability of assessment scores is a product of the difference in land cover distributions and the percentage of canopy cover within the population density classes and mapping zones (Fig. OH-10; Tables OH-11 through 14).




Figure OH-11.—Planting priority index for county subdivisions. The higher the index value, the greater priority for planting.


Priority Areas for Tree Planting

Priority areas for planting tend to be highest in more urbanized areas due to higher population density (Fig. OH-11; Tables OH-15 through 17). These index values can also be produced using high resolution cover data to determine local planting priority areas (e.g., neighborhoods).

Urban Tree Benefits

The following forest attributes are estimated for the urban or community land in Ohio (Table OH-1). These are rough estimates of values. More localized data are needed for more precise estimates, but these values reveal first-order approximations.

- 133.5 million trees
- 25.5 million metric tons of C stored (\$581.4 million value)
- 840,000 metric tons/year of C sequestered (\$19.2 million value)
- 21,930 metric tons/year total pollution removal (\$178.1 million value)
 - 311 metric tons/year of CO removed (\$438,000 value)
 - 3,832 metric tons/year NO₂ removed (\$38.0 million value)
 - 9,157 metric tons/year of O₃ removed (\$90.7 million value)
 - 1,934 metric tons/year of SO₂ removed (\$4.7 million value)
 - 6,693 metric tons/year of PM₁₀ removed (\$44.3 million value)




Summary

The data presented in this report provide a better understanding of Ohio's urban and community forests. This information can be used to advance urban and community forest policy and management that could improve environmental quality and human health throughout the State.

These data establish a baseline to assess future change and can be used to understand:

- Extent of the urban and community forest resource
- Variations in the resource across the State
- Magnitude and value of the urban and community forest resource
- Urban growth in Ohio
- Implications of policy decisions related to urban sprawl and urban and community forest management

WISCONSIN'S URBAN AND COMMUNITY FORESTS



Statewide Summary

Urban or community land in Wisconsin comprises about 5.6 percent of the state land area in 2000, an increase from 4.9 percent in 1990. Statewide tree canopy cover averages 38.9 percent and tree cover in urban or community areas is about 19.8 percent, with 16.2 percent impervious surface cover and 23.6 percent of the total green space covered by tree canopy cover. Statewide, urban or community land in Wisconsin has an estimated 74 million trees, which store about 14.1 million metric tons of carbon (\$321.5 million), and annually remove about 466,000 metric tons of carbon (\$10.6 million) and 9,610 metric tons of air pollution (\$82 million) (Table WI-1).

Tables WI-2 through WI-17 are not printed in this report but are available on the CD located on the inside back cover and at <http://nrs.fs.fed.us/data/urban>.

Table WI-1.—Statewide summary of population, area, population density, tree canopy and impervious surface land cover, and urban tree benefits in urban, community, and urban or community areas.

Wisconsin		Statewide	Urban ^a	Community ^b	Urban or Community ^c	
Population	2000	5,363,675	3,663,643	3,789,670	n/a	
	1990	4,891,769	3,211,956	3,466,580	n/a	
	% Change (1990-2000)	9.6	14.1	9.3	n/a	
	% Total population (2000)	100.0	68.3	70.7	n/a	
Total area	km ² (2000)	169,638.6	4,296.8	7,333.6	8,365.9	
	km ² (1990)	169,638.6	3,649.1	6,459.1	7,331.1	
	% Change (1990-2000)	0.0	17.7	13.5	14.1	
Land area	km ² (2000)	140,236.7	4,197.0	6,843.7	7,849.8	
	% Land area (2000)	100.0	3.0	4.9	5.6	
	km ² (1990)	140,236.7	3,565.4	6,034.8	6,878.5	
	% Land area (1990)	100.0	2.5	4.3	4.9	
	% Change (1990-2000)	0.0	17.7	13.4	14.1	
Population density (people/land area km ²)	2000	38.2	872.9	553.7	n/a	
	1990	34.9	900.9	574.4	n/a	
	% Change (1990-2000)	9.6	-3.1	-3.6	n/a	
Tree canopy cover (2000)	km ²	54,572.7	550.0	1,385.3	1,552.7	
	% Land area	38.9	13.1	20.2	19.8	
	Per capita (m ² /person)	10,174.5	150.1	365.5	n/a	
	% Canopy green space ^d	39.5	17.3	24.4	23.6	
Total green space (2000) ^e	km ²	138,140.0	3,178.8	5,666.4	6,575.3	
	% Land area	98.5	75.7	82.8	83.8	
Available green space (2000) ^f	km ²	83,567.8	2,628.9	4,281.3	5,022.8	
	% Land area	59.6	62.6	62.6	64.0	
Impervious surface cover (2000)	km ²	2,096.8	1,018.2	1,177.3	1,274.5	
	% Land area	1.5	24.3	17.2	16.2	
	Per capita (m ² /person)	390.9	277.9	310.7	n/a	
Urban tree benefits (2000)	Estimated number of trees	n/a	26,200,000	66,100,000	74,000,000	
	Carbon					
	Carbon stored (metric tons)	n/a	5,000,000	12,600,000	14,100,000	
	Carbon stored (\$)	n/a	\$114,000,000	\$287,300,000	\$321,500,000	
	Carbon sequestered (metric tons/year)	n/a	165,000	416,000	466,000	
	Carbon sequestered (\$/year)	n/a	\$3,762,000	\$9,485,000	\$10,625,000	
	Pollution					
	CO removed (metric tons/year)	n/a	56	142	159	
	CO removed (\$/year)	n/a	\$79,100	\$199,200	\$223,300	
	NO ₂ removed (metric tons/year)	n/a	610	1,536	1,721	
	NO ₂ removed (\$/year)	n/a	\$6,040,300	\$15,212,700	\$17,051,200	
	O ₃ removed (metric tons/year)	n/a	1,670	4,205	4,713	
	O ₃ removed (\$/year)	n/a	\$16,540,000	\$41,656,000	\$46,690,000	
	SO ₂ removed (metric tons/year)	n/a	164	414	464	
	SO ₂ removed (\$/year)	n/a	\$398,300	\$1,003,200	\$1,124,400	
PM ₁₀ removed (metric tons/year)	n/a	904	2,276	2,551		
PM ₁₀ removed (\$/year)	n/a	\$5,977,900	\$15,055,500	\$16,875,000		
Total pollution removal (metric tons/year)	n/a	3,400	8,570	9,610		
Total pollution removal (\$/year)	n/a	\$29,000,000	\$73,100,000	\$82,000,000		

^a Urban land is based on population density and was delimited using the United States Census definitions of urbanized areas and urban clusters. ^b Community land is based on jurisdictional or political boundaries of communities based on United States Census definitions of incorporated or census designated places. ^c Urban or communities is land that is urban, community, or both. Communities may include all, some, or no urban land within their boundaries. ^d Canopy green space is the tree canopy cover divided by total green space. ^e Total green space (TGS) is total area – impervious surface cover – water. ^f Available green space (AGS) is total green space – tree canopy cover (if the calculated value is less than 0, then value set at 0).




Figure WI-1.—Urban or community land in 2000; urban area relative to community boundaries.




Figure WI-2.—2000 population within county subdivision boundaries.

Human Population Characteristics and Trends

The population in Wisconsin increased 9.6 percent, from 4,891,769 in 1990 to 5,363,675 in 2000 (Table WI-1). In Wisconsin, 68.3 percent of the State’s population is in urban areas (Fig. WI-1), and 70.7 percent of the population is within communities (Fig. WI-2).

Urban and Community Land

Urban land comprises 3.0 percent of the land area of Wisconsin, while lands within communities make up 4.9 percent of the State (Fig. WI-1). Between 1990 and 2000, urban area increased 17.7 percent, while community land increased from 4.3 to 4.9 percent (Table WI-1). Urban area in Wisconsin is projected to increase to 8.3 percent by 2050, based on average urban growth pattern of the 1990s (Nowak and Walton 2005). Both urban land (attaining minimum population density) and community land (political boundaries) increased from 1990 to 2000. The percentages are calculated using the total (water and land) area of the geopolitical units derived from U.S. Census cartographic boundary data. Percent urban land varied across the State (Fig. WI-3; Tables WI-2 through 4).




Figure WI-3.—Percent of county subdivision area classified as urban land in 2000.




Figure WI-4.—Percentage tree canopy cover.




Figure WI-5.—Percentage tree canopy cover within county subdivisions.

Tree Canopy Cover Characteristics

Tree canopy cover in Wisconsin averages 38.9 percent (Fig. WI-4), with 98.5 percent total green space, 39.5 percent canopy green space, and 10,174.5 m² of canopy cover per capita. Average tree cover in urban areas in Wisconsin was 13.1 percent, with 75.7 percent total green space, 17.3 percent canopy green space, and 150.1 m² of canopy cover per capita. Within community lands in Wisconsin, average tree cover was 20.2 percent, with 82.8 percent total green space, 24.4 percent canopy green space, and 365.5 m² of canopy cover per capita (Table WI-1). Tree canopy cover, canopy green space, and tree cover per capita varied among communities, county subdivisions, and counties (Fig. WI-5 through 6; Tables WI-5 through 7).




Figure WI-6.—Percentage tree canopy green space in county subdivisions.




Figure WI-7.—Percentage impervious surface cover.




Figure WI-8.—Percentage impervious surface cover within county subdivisions.

Impervious Surface Cover Characteristics

Average impervious surface cover in Wisconsin is 1.5 percent of the land area (Fig. WI-7), with 390.9 m² of impervious surface cover per capita. Average impervious surface cover in urban areas was 24.3 percent, with 277.9 m² of impervious surface cover per capita. Within community lands in Wisconsin, average impervious surface cover was 17.2 percent with 310.7 m² of impervious surface cover per capita (Table WI-1). Impervious surface cover varied across the state (Fig. WI-8; Tables WI-5 through 7).




Figure WI-9.—Classified land cover.

Figure WI-10.—Relative comparisons of tree cover for county subdivisions.

Classified Land-cover Characteristics

Wisconsin's land cover is dominated by agricultural land (Fig. WI-9). The characteristics as a percent of the total land area in Wisconsin are (Tables WI-8 through 10):

- Agricultural – 47.2 percent
- Forested – 39.7 percent
- Developed – 6.9 percent
- Wetland – 3.4 percent
- Grassland – 1.8 percent
- Scrub/Shrub – 1.1 percent
- Barren – 0.1 percent

Relative Comparisons of Tree Cover

Out of the 630 Wisconsin communities, 16 received a rating of excellent and 353 received a rating of poor (Table WI-12). Of the 1,901 county subdivisions, 51 had a rating of excellent and 976 were rated poor (Fig. WI-10, Table WI-13); and out of 72 counties, seven were given a rating of excellent and 22 were given a rating of poor (Table WI-14). Variability of assessment scores is a product of the difference in land cover distributions and the percentage of canopy cover within the population density classes and mapping zones (Fig. WI-10; Tables WI-11 through 14).




Figure WI-11.—Planting priority index for county subdivisions. The higher the index value, the greater priority for planting.


Priority Areas for Tree Planting

Priority areas for planting tend to be highest in more urbanized areas due to higher population density (Fig. WI-11; Tables WI-15 through 17). These index values can also be produced using high resolution cover data to determine local planting priority areas (e.g., neighborhoods).

Urban Tree Benefits

The following forest attributes are estimated for the urban or community land in Wisconsin (Table WI-1). These are rough estimates of values. More localized data are needed for more precise estimates, but these values reveal first-order approximations.

- 74 million trees
- 14.1 million metric tons of C stored (\$321.5 million value)
- 466,000 metric tons/year of C sequestered (\$10.6 million value)
- 9,610 metric tons/year total pollution removal (\$82 million value)
 - 159 metric tons/year of CO removed (\$223,300 value)
 - 1,721 metric tons/year NO₂ removed (\$17.1 million value)
 - 4,713 metric tons/year of O₃ removed (\$46.7 million value)
 - 464 metric tons/year of SO₂ removed (\$1.1 million value)
 - 2,551 metric tons/year of PM₁₀ removed (\$16.9 million value)



Summary

The data presented in this report provide a better understanding of Wisconsin's urban and community forests. This information can be used to advance urban and community forest policy and management that could improve environmental quality and human health throughout the State.

These data establish a baseline to assess future change and can be used to understand:

- Extent of the urban and community forest resource
- Variations in the resource across the State
- Magnitude and value of the urban and community forest resource
- Urban growth in Wisconsin
- Implications of policy decisions related to urban sprawl and urban and community forest management

LITERATURE CITED

- Dwyer, J.F.; Nowak, D.J.; Noble, H.M.; Sisinni, S.M. 2000. **Assessing our nation's urban forests: connecting people with ecosystems in the 21st century.** Gen. Tech. Rep. PNW-490. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 540 p.
- Fankhauser, S. 1994. **The social costs of greenhouse gas emissions: an expected value approach.** The Energy Journal. 15(2): 157-184.
- Greenfield, E.J.; Nowak D.J.; Walton, J.T. 2009. **Assessment of 2001 NLCD percent tree and impervious cover estimates.** Photogrammetric Engineering and Remote Sensing. 75(11): 1279-1286.
- Homer, C.; Huang, C.; Yang, L.; Wylie, B.; Coan, M. 2004. **Development of a 2001 national land cover database for the United States.** Photogrammetric Engineering and Remote Sensing. 70(7): 829-840.
- Homer, C.G.; Gallant, A. 2001. **Partitioning the conterminous United States into mapping zones for Landsat TM land cover mapping.** Unpublished U.S. Geologic Survey report. <http://landcover.usgs.gov/pdf/homer.pdf>. (1 August 2008).
- Homer, C.; Dewitz, J.; Fry, J.; Coan, M.; Hossain, N.; Larson, C.; Herold, N.; McKerrow, A.; VanDriel, J.N.; Wickham, J. 2007. **Completion of the 2001 national land cover database for the coterminous United States.** Photogrammetric Engineering and Remote Sensing. 73(4): 337-341.
- Murray, F.J.; Marsh, L.; Bradford, P.A. 1994. **New York state energy plan Vol. II: issue reports.** Albany, NY: New York State Energy Research and Development Authority.
- National Climatic Data Center. 2000. **Integrated surface hourly observations 2000.** Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration. [CD-ROM].
- Nowak, D.J. 1993. **Compensatory value of an urban forest: an application of tree-value formula.** Journal of Arboriculture. 19(3): 173-177.
- Nowak, D.J. 1994a. **Atmospheric carbon dioxide reduction by Chicago's urban forest.** In: McPherson, E.G; Nowak, D.J.; Rowntree, R.A. Chicago's urban forest ecosystem: results of the Chicago urban forest climate project. Gen. Tech. Rep. NE-186. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station: 83-94.
- Nowak, D.J. 1994b. **Urban forest structure: the state of Chicago's urban forest.** In: McPherson, E.G; Nowak, D.J.; Rowntree, R.A. Chicago's urban forest ecosystem: results of the Chicago urban forest climate project. Gen. Tech. Rep. NE-186. Radnor, PA: U.S. Department of Agriculture, Forest Service Northeastern Research Station: 3-18 and 140-164 [appendix].

- Nowak, D.J.; Crane, D.E. 2000. **The Urban Forest Effects (UFORE) model: quantifying urban forest structure and functions.** In: Hansen, M.; Burk, T., eds. Integrated tools for natural resources inventories in the 21st century, proceedings of the IUFRO conference; 1998 August 16-20; Boise, ID. Gen. Tech. Rep. NC-212. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station: 714-720.
- Nowak, D.J.; Crane, D.E. 2002. **Carbon storage and sequestration by urban trees in the United States.** Environmental Pollution. 116(3): 381-389.
- Nowak, D.J.; Crane, D.E.; Stevens, J.C. 2001a. **Syracuse's urban forest resource.** In: Nowak, D.J.; O'Connor, P., comps. Syracuse urban forest master plan: guiding the city's forest resource in the 21st century. Gen. Tech. Rep. NE-287. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station: 9-14.
- Nowak, D.J.; Crane, D.E.; Stevens, J.C. 2006d. **Air pollution removal by urban trees and shrubs in the United States.** Urban Forestry and Urban Greening. 4: 115-123.
- Nowak, D.J.; Hoehn, R.; Crane, D.E.; Stevens, J.C.; Walton, J.T. 2006a. **Assessing urban forest effects and values: Casper, WY's urban forest.** Res. Bull. NRS-4. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 20 p.
- Nowak, D.J.; Hoehn, R.; Crane, D.E.; Stevens, J.C.; Walton, J.T. 2006b. **Assessing urban forest effects and values: Minneapolis' urban forest.** Res. Bull. NE-166. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 20 p.
- Nowak, D.J.; Hoehn, R.; Crane, D.E.; Stevens, J.C.; Walton, J.T. 2006c. **Assessing urban forest effects and values: Washington D.C.'s urban forest.** Resour. Bull. NRS-1. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 24 p.
- Nowak, D.J.; Hoehn, R.; Crane, D.E.; Stevens, J.C.; Walton, J.T. 2007a. **Assessing urban forest effects and values: New York's urban forest.** Resour. Bull. NRS-9. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 22 p.
- Nowak, D.J.; Hoehn, R.; Crane, D.E.; Stevens, J.C.; Walton, J.T. 2007b. **Assessing urban forest effects and values: Philadelphia's urban forest.** Resour. Bull. NRS-7. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 22 p.
- Nowak, D.J.; Hoehn, R.; Crane, D.E.; Stevens, J.C.; Walton, J.T. 2007c. **Assessing urban forest effects and values: San Francisco's urban forest.** Resour. Bull. NRS-8. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 22 p.

- Nowak, D.J.; Noble, M.H.; Sisinni, S.M.; Dwyer, J.F. 2001b. **People and trees: assessing the U.S. urban forest resource.** *Journal of Forestry*. 99(3): 37-42.
- Nowak, D.J.; Walton, J.T.; Dwyer, J.F.; Kaya, L.G.; Myeong, S. 2005. **The increasing influence of urban environments on U.S. forest management.** *Journal of Forestry*. 103(8): 377-382.
- Nowak, D.J.; Walton, J.T. 2005. **Projected urban growth and its estimated impact on the U.S. forest resource (2000-2050).** *Journal of Forestry*. 103(8): 383-389.
- Ottinger, R.L.; Wooley, D.R.; Robinson, N.A.; Hodas, D.R.; Babb, S.E.; Buchanan, S.C.; Chernick, P.L.; Caverhill, E.; Krupnick, A.; Fritsche, U. 1990. **Environmental costs of electricity.** White Plains, NY: Oceana Publications. 769 p.
- U.S. Census Bureau. n.d. **www.census.gov.** (January 2007).
- U.S. Department of Interior, Geologic Survey. 2008. **Multi-resolution land characteristics consortium.** **www.mrlc.gov.** (1 August 2008).
- U.S. Department of Labor Bureau of Labor Statistics. n.d. **www.bls.gov/ppi/** (June 2007).
- U.S. Environmental Protection Agency. 2003. **National air quality and emissions trends report: 2003 special studies edition.** Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Emissions Monitoring and Analysis Division.
- U.S. Environmental Protection Agency. n.d. **www.epa.gov/oar/data** (June 2008).
- Walton, J.T. 2005. **An investigation of national tree canopy assessments applied to urban forestry.** Syracuse, NY: State University of New York, College of Environmental Science and Forestry. 95 p. Ph.D. dissertation.
- Yang, L.; Huang, C.; Homer, C.G.; Wylie, B.K.; Coan, M.J. 2003. **An approach for mapping large-area impervious surfaces: synergistic use of Landsat-7 ETM+ and high spatial resolution imagery.** *Canadian Journal of Remote Sensing*. 29(2): 230-240.
- Zhu, Z. 1994. **Forest density mapping in the lower 48 states: a regression procedure.** Res. Pap. SO-280. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Research Station. 11 p.

APPENDIX

Urban Forest Data: States of the North Central East Region

The following tables are generated to support state reports on urban and community forests of the North Central East states of Illinois, Indiana, Michigan, Ohio, and Wisconsin. For specific state data tables use the CD accompanying this publication and search within the regional or state folder, or go to: <http://www.nrs.fs.fed.us/data/urban>.

State Specific Tables:

Table 1.—Statewide summary of population, area, population density, tree canopy and impervious surface land cover, and urban tree benefits in urban, community, and urban or community areas.

Table 2.—2000 population characteristics, population change (1990-2000), and percent of land classified as urban within communities.

Table 3.—2000 population characteristics, population change (1990-2000), percent of land classified as urban or as communities within county subdivisions.

Table 4.—2000 population characteristics, population change (1990-2000), percent of land classified as urban or as communities within counties.

Table 5.—Tree canopy and impervious surface cover characteristics by community.

Table 6.—Tree canopy and impervious surface cover characteristics by county subdivision.

Table 7.—Tree canopy and impervious surface cover characteristics by county.

Table 8.—Land area, tree canopy cover, and available green space distributed within generalized land cover categories for communities.

Table 9.—Land area, tree canopy cover, and available green space distributed within generalized land cover categories for county subdivisions.

Table 10.—Land area, tree canopy cover, and available green space distributed within generalized land cover categories for counties.

Table 11.—Statistical summary of mapping zone values used to calculate urban and community forestry assessment.

Table 12.—Urban and community forestry assessment by community.

Table 13.—Urban and community forestry assessment by county subdivisions.

Table 14.—Urban and community forestry assessment by counties.

Table 15.—Planting priority index for communities.

Table 16.—Planting priority index for county subdivisions.

Table 17.—Planting priority index for counties.

Nowak, David J.; Greenfield, Eric J. 2010. **Urban and community forests of the North Central East region: Illinois, Indiana, Michigan, Ohio, Wisconsin.** Gen. Tech. Rep. NRS-54. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 54 p.

This report details how land cover and urbanization vary within the states of Illinois, Indiana, Michigan, Ohio, and Wisconsin by community (incorporated and census designated places), county subdivision, and county. Specifically this report provides critical urban and community forestry information for each state including human population characteristics and trends, changes in urban and community lands, tree canopy and impervious surface cover characteristics, distribution of land-cover classes, a relative comparison of urban and community forests among local government types, determination of priority areas for tree planting, and a summary of urban tree benefits. Report information can improve the understanding, management, and planning of urban and community forests. This data is reported for each state on the CD provided in the back of this book and may be accessed by state at: <http://www.nrs.fs.fed.us/data/urban>.

KEY WORDS: urban forestry, tree cover, impervious cover, classified land cover, ecosystem services, urbanization

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and, where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternate means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue S.W., Washington, DC 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.



Capitalizing on the strengths of existing science capacity in the Northeast and Midwest to attain a more integrated, cohesive, landscape-scale research program

www.nrs.fs.fed.us